

A very large glitch in PSR B2334+61

J. P. Yuan^{1,2}, R. N. Manchester³, N. Wang^{1,*}, X. Zhou¹, Z. Y. Liu¹ and Z. F. Gao^{1,2}

¹*Urumqi Observatory, NAOC, 40-5 South Beijing Road, Urumqi, Xinjiang, China, 830011*

²*Graduate University of CAS, 19A Yuquan road, Beijing, China, 100049*

³*CSIRO Astronomy and Space Science, Australia Telescope National Facility, PO Box 76, Epping, NSW 1710, Australia*

**E-mail: na.wang@uao.ac.cn*

ABSTRACT

Seven years of pulse time-of-arrival measurements have been collected from observations of the young pulsar PSR B2334+61 using the Nanshan radio telescope of Urumqi Observatory. A phase-connected solution has been obtained over the whole data span, 2002 August to 2009 August. This includes a very large glitch that occurred between 2005 August 26 and September 8 (MJDs 53608 and 53621). The relative increase in rotational frequency for this glitch, $\Delta\nu_g/\nu \sim 20.5 \times 10^{-6}$, is the largest ever seen. Although accounting for less than 1% of the glitch, there were two well-defined exponential decay terms with time constants of 21 and 147 days respectively. There was also a large long-term increase in the spindown rate with $\Delta\dot{\nu}_p/\dot{\nu} \sim 0.011$ at the time of the glitch. A highly significant oscillation with a period of close to one year is seen in the post-glitch residuals. It is very unlikely that this can be accounted for by a pulsar position error or proper motion – it appears to result from effects interior to the neutron star. Implications of these results for pulsar glitch models are discussed.

Subject headings: stars: neutron — pulsars: general — pulsars: individual (PSR B2334+61)

1. Introduction

Pulsars are well known for their very stable spin periods. However, long-term timing observations of young pulsars have revealed two categories of rotational irregularities: timing noise which is usually characterized by continuous and erratic fluctuation in rotation period, and glitches which are sudden increases in their rotation rates. Typical increases in rotational

frequency during a glitch are of the order of $\Delta\nu_g/\nu \sim 10^{-9}$ to 10^{-6} . Large glitches are predominately found in young pulsars with characteristic ages between 10 kyr and 1000 kyr and are often followed by a relaxation process in which some of the glitch decays. It is believed that glitches and post-glitch variations reflect the dynamics of the interior of the neutron star rather than magnetospheric phenomena. Observing glitches and measuring their subsequent decay processes, in addition to providing insights into the phenomenon itself, offer one of the few probes of the neutron star structure and thus the physics of ultra-dense matter (e.g., Larson & Link 2002; Alpar & Baykal 2006).

The young pulsar PSR B2334+61 (PSR J2337+6151) was discovered in the Princeton-NRAO survey using the 92-m radio telescope at Green Bank (Dewey et al. 1985); it is located near the center of, and probably associated with, the supernova remnant G114.3+0.3 (Fürst et al. 1993). The pulsar has a rotational period $P \sim 0.495$ s and a period derivative $\dot{P} \sim 1.9 \times 10^{-13}$, implying a characteristic age $\tau_c = P/(2\dot{P}) \sim 41$ kyr. B2334+61’s spin-down power, $\dot{E} = -4\pi^2 I \nu \dot{\nu}$, is 6.23×10^{34} erg s $^{-1}$, where the moment of inertia I is taken to be 10^{45} g cm ^{2} . This suggests that the pulsar may be a γ -ray emitter, although its distance of ~ 3 kpc puts it at the bottom end of the $\dot{E}^{1/2} d^{-2}$ metric for pulsars detectable by the *Fermi* Space Telescope (Abdo et al. 2010). X-ray emission from the pulsar was first detected in a ROSAT pointing (Becker et al. 1996) and was shown to be thermal using an *XMM-Newton* observation (McGowan et al. 2006).

In this Letter we present the timing solution of PSR B2334+61 obtained from seven years of observation with the Nanshan 25-m telescope. The timing analysis reveals the largest known glitch ever observed, with a fractional frequency increase of $\Delta\nu_g/\nu \sim 20.5 \times 10^{-6}$, a double exponential decay of a small part of the glitch, a large permanent increase in spindown rate and a quasi-sinusoidal oscillation apparently triggered by the glitch. We give a short discussion about the implications of these results for glitch theories.

2. OBSERVATION AND ANALYSIS

Timing observations of PSR B2334+61 using the 25-m Nanshan radio telescope at Urumqi Observatory commenced in 2002 July. The observing system is described in Wang et al. (2001). In brief, the two hands of circular polarization at a frequency of 1540 MHz are fed through a $2 \times 128 \times 2.5$ MHz-channel filter bank and digitized at 1 ms intervals. The observation time for this pulsar was 16 minutes.

Off-line data reduction was performed in stages using the PSRCHIVE package¹. The data were dedispersed and summed to produce a total intensity profile. Local times-of-arrival (TOAs) were determined by correlating this profile with a standard pulse profile of high signal-to-noise ratio. Correction of TOAs to the Solar-System barycentre and fitting of the timing model were done using TEMPO2² with the Jet Propulsion Laboratory DE405 ephemeris (Standish 1998). TOAs were weighted by the inverse-square of their estimated uncertainty. The basic timing model is the Taylor series

$$\phi(t) = \phi_0 + \nu(t - t_0) + \frac{1}{2}\dot{\nu}(t - t_0)^2 + \frac{1}{6}\ddot{\nu}(t - t_0)^3 + \frac{1}{24}\dddot{\nu}(t - t_0)^4 \quad (1)$$

where ϕ_0 is the pulsar phase at the reference barycentric time t_0 . Glitches can be modelled as combinations of changes in ν , $\dot{\nu}$ and $\ddot{\nu}$, of which parts may recover on various timescales:

$$\nu(t) = \nu_0(t) + \Delta\nu_p + \Delta\dot{\nu}_p t + \frac{1}{2}\Delta\ddot{\nu}_p t^2 + \Delta\nu_d e^{-t/\tau_d} \quad (2)$$

$$\dot{\nu}(t) = \dot{\nu}_0(t) + \Delta\dot{\nu}_p + \Delta\ddot{\nu}_p t + \Delta\dot{\nu}_d e^{-t/\tau_d} \quad (3)$$

$$\ddot{\nu}(t) = \ddot{\nu}_0(t) + \Delta\ddot{\nu}_p + \Delta\ddot{\nu}_d e^{-t/\tau_d} \quad (4)$$

where $\Delta\nu_p$ and $\Delta\dot{\nu}_p$ are permanent changes in ν and $\dot{\nu}$ relative to the pre-glitch solution, $\Delta\nu_d$ is the amplitude of a decaying component with a time constant of τ_d , $\Delta\dot{\nu} = -\Delta\nu_d/\tau_d$ and $\Delta\ddot{\nu}_d = \Delta\nu_d/\tau_d^2$. The total frequency jump at the time of the glitch is $\Delta\nu_g = \Delta\nu_p + \Delta\nu_d$. The degree of recovery is often described by the parameter $Q = \Delta\nu_d/\Delta\nu_g$. If required, multiple decay terms with different decay time constants can be fitted; the total frequency jump is then the sum of the permanent and decaying frequency jumps, and similarly for $\dot{\nu}$ and $\ddot{\nu}$.

3. RESULTS

The data presented here were recorded between 2002 July 31 and 2009 August 1 (MJDs 52486 and 55044). Figure 1 gives an overview of the results of a timing analysis, showing that a large glitch with $\Delta\nu_g \approx 41 \times 10^{-6}$ Hz occurred between 2005 August 26 and September 7 (MJDs 53608 and 53621). The regular timing observations of this pulsar at Nanshan allow us to track the recovery process in detail. Most of the frequency jump $\Delta\nu_g$ persists beyond the end of the data span, but the expanded $\Delta\nu$ plot given in Figure 1(b) and the plot of $\dot{\nu}$ given in Figure 1(c) show that there was an initial exponential decay with a timescale of

¹See <http://psrchive.sourceforge.net>

²See <http://www.atnf.csiro.au/research/pulsar/tempo2>

~ 150 days and an apparently permanent increase in slow-down rate $|\dot{\nu}|$. As is described in detail below, there was an additional more rapid exponential decay with a timescale of ~ 21 days following the glitch.

The rotational and positional parameters for PSR B2334+61 from fitting of the timing model, Equation 1, to the pre- and post-glitch data are given in Table 1. Pulsar frequencies are in TDB units and uncertainties given in parentheses are in the last quoted digit and are twice the values given by TEMPO2. The position was derived from the pre-glitch data assuming the proper motion given by Hobbs et al. (2004) and held fixed for the post-glitch solution. To avoid most of the short-term post-glitch decay, the post-glitch solution was fitted to data starting approximately one year after the glitch. Even with this, there were significant systematic period variations beyond the second-derivative term, most of which were absorbed by a third-derivative term.

Figure 1(c) shows that there was a very significant long-term increase in $|\dot{\nu}|$ at the time of the glitch; the values of $\dot{\nu}$ given in Table 1 reflect this change. Furthermore, there was a significant long-term change in $\ddot{\nu}$ at the time of the glitch, with the rate of decrease of $|\dot{\nu}|$ increasing by more than four after the glitch. The corresponding braking indices are 10.5 ± 0.2 and 46.8 ± 0.3 before and after the glitch, respectively.

Table 2 gives glitch parameters from a fitting of a model to the timing data. Because of the large amplitude of the glitch, the final fit was obtained in several stages. For the first stage, fits the data span were terminated at MJD 54000, just over a year after the glitch, to avoid contamination by the long-term decay in $\dot{\nu}$. $\Delta\dot{\nu}_p$ was held fixed at the value given by subtraction of the pre-glitch value of $\dot{\nu}$ given in Table 1 from the post-glitch value, viz, $-7.625 \times 10^{-15} \text{ s}^{-2}$. Then, based on Figure 1, a single decay with an assumed time constant of 150 d was fitted to the TOAs, giving values of $\Delta\nu_p \sim 4.1203 \times 10^{-5} \text{ Hz}$ and $\Delta\nu_{d1} \sim 1.30 \times 10^{-7} \text{ Hz}$. Post-fit residuals from this fit are shown in the upper part of Figure 2. It is clear from this plot that there was an additional initial decay with time constant ~ 20 d. Fitting for this with the parameters of the longer decay held fixed gave $\Delta\nu_{d2} \sim 1.9 \times 10^{-7} \text{ Hz}$ and $\tau_{d2} = 18.5 \text{ d}$. Residuals for this fit shown in the lower part of Figure 2 demonstrate that the observed TOAs over this data span are very well modelled by increments in ν and $\dot{\nu}$ together with two exponential decays with time constants around 20 d and 150 d respectively.

Using these parameters as starting points, it was then possible to fit the entire dataset from 2002 to 2009 to give the glitch parameters listed in Table 2. The pulsar position, proper motion, ν , $\dot{\nu}$ and $\ddot{\nu}$ values were held fixed at the pre-glitch values from Table 1. All other parameters were solved for with the fit being iterated several times to ensure

convergence³. There was a gap of 12 days between observations around the glitch; the assumed glitch epoch is in the middle of this gap. Although most derived parameters are essentially independent of the assumed epoch, the amplitudes of the decaying components do change significantly. Quoted uncertainties in the glitch parameters include a contribution from the uncertainty in the glitch epoch. Residuals from this final fit are shown in Figure 3. Values for $\Delta\dot{\nu}_p$ and $\Delta\ddot{\nu}_p$ are close to the differences between $\dot{\nu}$ and $\ddot{\nu}$ for the post- and pre-glitch fits (Table 1) as expected. They do not coincide exactly (within the estimated errors) mainly because the exponential terms included in the final fit absorb some of these terms.

This final fit clearly represents the glitch and the subsequent decay extremely well. To put the scale of the residuals in Figure 3 into context, $\Delta\nu_g$ represents a phase gradient of one turn every 6.2 hours and the $\Delta\dot{\nu}_p$ and $\Delta\ddot{\nu}_p$ terms correspond to a total phase contribution over the post-glitch data span of 66.2 and 2.76 turns respectively. The final rms residual corresponds to 0.0035 turns. Although good, the final fit is not perfect, with a reduced χ^2 of 3.32. The χ^2 is dominated by systematic deviations from the fit after the glitch. These appear to be quasi-sinusoidal; a Lomb-Scargle spectral analysis shows a periodicity in the post-glitch post-fit residuals of 364 d with an uncertainty of ~ 5 d and a significance (false-alarm probability) of 0.0005. The fact that the period of the post-glitch oscillations is equal to a year within the uncertainty immediately suggests that the oscillations are due to a position error or proper motion. However, we believe that this is not the correct explanation.

The position derived from the pre-glitch data (Table 1) is fully consistent with the position extrapolated to MJD 53100 based on the position and proper motion obtained by Hobbs et al. (2004) using Jodrell Bank data spanning 16 years: $23^{\text{h}} 37^{\text{m}} 05.779(28)$, $+61^{\circ}51'01''.54(17)$. A fit of position only to the post-glitch data, holding all glitch and pulse frequency parameters constant at the final-fit values, moves the position by $+0.059$ of right ascension and $+0.36$ of declination, about four times the combined uncertainties. Compared to the extrapolated Hobbs et al. (2004) position, the shifts are about twice the combined uncertainties. Including position, proper motion, ν , $\dot{\nu}$, $\ddot{\nu}$ and all glitch terms in a fit to the whole data set produced small changes in all glitch parameters (less than the uncertainties given in Table 2) and gave a proper motion of $+111(24)$ mas yr^{-1} in the right ascension direction and $+75(23)$ mas yr^{-1} in declination. These values are inconsistent with the Hobbs et al. (2004) values. Furthermore, the final post-fit residuals have significant annual terms of opposite sign before and after the glitch. These considerations strongly suggest that the solution illustrated in Figure 3 is the best possible, that the oscillations are confined to

³Solving for the pulsar position, ν , $\dot{\nu}$ and $\ddot{\nu}$ also converged well and did not change any of the parameters outside their uncertainty range, demonstrating that the covariance between the various parameters is small.

post-glitch epochs and that they are not caused by a position error or proper motion of the pulsar.

There is no evidence for a change in pulse shape at the time of the glitch, but the limits are rather poor as the typical signal-to-noise ratio from each 16-minute observation was only about 10.

4. DISCUSSION

The glitch in PSR B2334+61 is the largest in $\Delta\nu_g/\nu$ terms (and equal largest in $\Delta\nu_g$ terms) ever observed for any radio pulsar and probably in any pulsar (including magnetars).⁴ With a characteristic age of 41 kyr it is in the age range of pulsars known to exhibit large glitches (see, e.g., Yuan et al. 2010). In fact, it lies very close in the $P - \dot{P}$ plane to PSR J1806–2125 which in 1998 had a glitch with $\Delta\nu_g/\nu \sim 16 \times 10^{-6}$ (Hobbs et al. 2002). There are other six pulsars close to PSR B2334+61 in the $P - \dot{P}$ plane: PSRs J1551–5310, J1737–3137, B1758–23, J1838–0453, J1841–0524 and B2000+32. For PSR B1758–23, six glitches of moderate size ($\Delta\nu_g/\nu$ in the range $(0.017 - 0.35) \times 10^{-6}$) have been detected. For PSRs J1737–3137 and J1841–0524 a single glitch of relative size $\sim 1.3 \times 10^{-6}$ and 1.0×10^{-6} respectively have been reported (Weltevrede et al. 2010). Other pulsars have not been reported to glitch so far. This is the first glitch reported for PSR B2334+61; since the pulsar was observed for 16 years from 1987 by Hobbs et al. (2004), the time between glitches is at least 22 years.

In contrast to the case for PSR J1806–2125, for PSR B2334+61 observations were obtained close to the date of the glitch allowing a detailed examination of the post-glitch behaviour. PSR B2334+61 continues the trend for minimal post-glitch recovery of $\Delta\nu_g$, i.e., small Q , for large glitches in middle-aged pulsars (Lyne et al. 2000; Yuan et al. 2010). Despite this, the good sampling of the post-glitch behaviour shows clear evidence for two distinct exponential recoveries and, together with permanent jumps in $\dot{\nu}$ and $\ddot{\nu}$, these provide a remarkably accurate description of the post-glitch behaviour. Previously, only the Vela pulsar has shown clear evidence for multiple exponential decays associated with a given glitch (Dodson et al. 2002). For Vela, the longest decay timescale observed by Dodson et al. (2002) is about 19 d, comparable to the shorter decay timescale observed for PSR B2334+61. There is evidence for longer exponential decays in the Vela pulsar glitches which ultimately become more linear (Lyne et al. 1996; Lyne et al. 2000; Wang et al. 2000).

⁴See the Glitch Table in the ATNF Pulsar Catalogue (www.atnf.csiro.au/research/pulsar/psrcat).

The standard model for large pulsar glitches envisages a sudden transfer of angular momentum from a superfluid component to the rest of the star – the vortex unpinning model (Anderson & Itoh 1975; Alpar et al. 1984). The theoretical understanding of glitches includes the trigger mechanism for the glitch, the strength of the vortex pinning and the post-glitch evolution. Larson & Link (2002) investigated a model in which vortex unpinning results from a sudden heating of the star, possibly due to a starquake or crustal movement, whereas Glampedakis & Andersson (2009) suggested that an r-mode instability may trigger a global unpinning of vortices leading to a glitch. Alpar et al. (1993) and earlier papers by these authors attribute the long-term post-glitch relaxation to vortex creep, with different regions of the star having different properties to account for the sometimes complicated post-glitch behaviour. Ruderman et al. (1998) suggested that crustal motions could account for essentially all glitch properties; see also Ruderman (2009).

The fractional jump in $\dot{\nu}$ at the time of the glitch was very large, about 15% (Table 2), much larger than the value of 1.7% found by Lyne et al. (2000) to represent most pulsars. For PSR B2334+61, $\Delta\dot{\nu}_g/\dot{\nu}$ is dominated by the fastest decay term, with the relative contributions for the 21-d decay, the 147-d decay and the permanent jump being 85%, 8% and 7%, respectively. In most glitch theories, $\Delta\dot{\nu}_g/\dot{\nu}$ represents the moment of inertia fraction which is in a superfluid state not tightly coupled to the neutron-star crust (see, e.g., Alpar et al. 1989, 1993). The large value for PSR B2334+61 suggests that a considerable fraction of the star is weakly coupled superfluid, with most of it returning to an equilibrium rotational state within about 20 days. About 1% of the rotational inertia appears to be more loosely coupled, leading to the 147-d decay, and a further 1% barely coupled at all.

The very significant quasi-permanent change in $\dot{\nu}$ associated with the PSR B2334+61 glitch (Figure 1) is unusual. Such clear quasi-permanent changes in $\dot{\nu}$ have previously only been observed in the Crab pulsar (Lyne et al. 1993). Although it is fairly common for only part of $\Delta\dot{\nu}_g$ to recover exponentially after a large glitch, in most cases there is a continuing approximately linear recovery of $\dot{\nu}$. As discussed above, this was seen in the Vela pulsar period, and other examples have been presented and discussed by Lyne et al. (2000) and Yuan et al. (2010). For PSR B2334+61, the fractional change $\Delta\dot{\nu}_p/\dot{\nu}$ is about 1.1%, about 25 times larger than the value of $\sim 4 \times 10^{-4}$ observed for the Crab pulsar. These sudden changes in $\dot{\nu}$ implies equally sudden changes in the effective braking torque at the time of the glitch since a change in the stellar moment of inertia of this magnitude is ruled out by the fact that $\Delta\nu_p/\nu$ is several orders of magnitude smaller. The observed change in $\dot{\nu}$ for PSR B2334+61 seems too large to be accounted for by exterior magnetic field changes as suggested for the Crab pulsar (Ruderman et al. 1998; Ruderman 2009). Furthermore, there was no major change in the pulse shape at the time of the glitch, suggesting that magnetospheric changes are not responsible. Therefore it is most likely that changes interior to the neutron star are

responsible. The effective braking torque is the difference between magnetospheric torques and internal torques believed to be due to vortex creep (e.g., Alpar et al. 1984). This model predicts a linear increase in $\dot{\nu}$ over long timescales as is seen in the Vela pulsar (Alpar et al. 1993). For PSR B2334+61 the observed value of $\dot{\nu}$ is two orders of magnitude smaller than for Vela, which is in accordance with the predictions of the vortex-creep model (Alpar & Baykal 2006).

The observed long-term step in $\ddot{\nu}$ (Table 2) could be interpreted as indicating a third exponential decay with a very long timescale. Large positive values of $\ddot{\nu}$ are commonly attributed to the long-term effect of (sometimes unseen) glitches (e.g., Hobbs et al. 2004; Alpar & Baykal 2006). For an exponential decay, $\Delta\ddot{\nu}_{d3} \sim \ddot{\nu}_p = \Delta\nu_{d3}/\tau_{d3}^2$. Since we do not know $Q_{d3} = \Delta\nu_{d3}/\Delta\nu_g$, we can just set a limit on $\tau_{d3} < 70$ years corresponding to $Q_{d3} = 1$. In principle, the observed post-glitch value of $\ddot{\nu}$ (Table 1) could be interpreted as $\Delta\ddot{\nu}_{d3}$ and used to solve for Q_{d3} and τ_{d3} , but it is inconsistent with this, implying a decay timescale $-\Delta\ddot{\nu}_{d3}/\Delta\ddot{\nu}_{d3}$ of less than one year. Evidently the shorter-term decays are not exactly exponential or other noise processes contribute to the observed value of $\ddot{\nu}$.

The oscillations in the post-glitch timing residuals are a remarkable feature of the timing behaviour of PSR B2334+61 (Figure 3). These oscillations have a period very close to one year, but in Section 3 we showed that an interpretation in terms of position error or proper motion is very improbable. Quasi-periodic oscillations in timing residuals have been seen in the Crab pulsar (Lyne et al. 1993) and many other pulsars (Hobbs et al. 2010). In none of these cases though are the oscillations clearly associated with a glitch although glitches have been mentioned as an excitation mechanism (e.g., Zhou et al. 2004; Timokhin 2007). The observed quasi-periodic oscillations have often been interpreted as due to Tkachenko oscillations in the superfluid vortex array (Ruderman 1970; Popov 2008). Under certain reasonable assumptions, these oscillations have a period given by

$$P_T \sim 1.77 R_6 P^{1/2} \text{yr} \quad (5)$$

where R_6 is the neutron-star radius in units of 10^6 cm and P is the pulsar period in seconds. For PSR B2334+61, this gives a period of $1.24 R_6$ yr, remarkably close to the observed period of 1 yr. This suggests that Tkachenko oscillations are a viable mechanism for the observed oscillations. Further investigation is needed to understand the triggering mechanism and whether or not the observed oscillation amplitude can be accounted for.

We thank George Hobbs for helpful discussions. This work is supported by National Basic Research Program of China – 973 Program 2009CB824800, NSFC projects 10673021, 10778631 and 10903019, Knowledge Innovation Program of The Chinese Academy Sciences

KJCX2-YW-T09, West Light Foundation of CAS (No. XBBS200920) and Xinjiang Natural Science Foundation (No. 2009211B35).

REFERENCES

- Abdo et al. 2010, ApJS, 187, 460
- Alpar, M. A., Anderson, P. W., Pines, D. & Shaham, J. 1984, ApJ, 276, 325
- Alpar, M. A., & Baykal, A. 2006, MNRAS, 372, 489
- Alpar, M. A., Chau, H. F., Cheng, K. S., & Pines, D. 1993, ApJ, 409, 345
- Alpar, M. A., Cheng, K. S., & Pines, D. 1989, ApJ, 346, 823
- Alpar, M. A., Langer, S. A., & Sauls, J. A. 1984, ApJ, 282, 533
- Anderson, P. W., & Itoh, N. 1975, Nature, 256, 25
- Becker W., Brazier K. T. S., Trümper J., 1996, A&A, 306, 464
- Dewey, R. J., Taylor, J. H., Weisberg, J. M., & Stokes, G. H. 1985, ApJ, 294, L25
- Dodson, R. G., McCulloch, P. M., & Lewis, D. R. 2002, ApJ, 564, L85
- Fürst, E., Reich, W., & Seiradakis, J. H. 1993, A&A, 276, 470
- Glampedakis, K., & Andersson, N. 2009, Phys. Rev. Lett., 102, 141101
- Hobbs, G., Lyne, A. G., & Kramer, M. 2010, MNRAS, 402, 1027
- Hobbs, G., Lyne, A. G., Kramer, M., Martin, C. E., & Jordan, C. 2004, MNRAS, 353, 1311
- Hobbs, G., et al. 2002, MNRAS, 333, L7
- Larson, M. B., & Link, B. 2002, MNRAS, 333, 613
- Lyne, A. G., Pritchard, R. S., Graham-Smith, F., & Camilo, F. 1996, Nature, 381, 497
- Lyne, A. G., Pritchard, R. S., & Smith, F. G. 1993, MNRAS, 265, 1003
- Lyne, A. G., Shemar, S. L., & Graham-Smith, F. 2000, MNRAS, 315, 534
- McGowan K. E., Zane S., Cropper M., Vestrand W. T., Ho C., 2006, ApJ, 639, 377

- Popov, S. B. 2008, *Astrophys. Space Sci.*, 317, 175
- Ruderman, M. 1970, *Nature*, 225, 619
- Ruderman, M. 2009, in *Astrophysics and Space Science Library*, Vol. 357, *Astrophysics and Space Science Library*, ed. W. Becker, 353–+
- Ruderman, M., Zhu, T., & Chen, K. 1998, *ApJ*, 492, 267
- Standish, E. M. 1998, *JPL Planetary and Lunar Ephemerides*, DE405/LE405, Memo IOM 312.F-98-048 (Pasadena: JPL), <http://ssd.jpl.nasa.gov/iau-comm4/de405iom/de405iom.pdf>
- Timokhin, A. N. 2007, *Astrophys. Space Sci.*, 308, 345
- Wang, N., Manchester, R. N., Pace, R., Bailes, M., Kaspi, V. M., Stappers, B. W., & Lyne, A. G. 2000, *MNRAS*, 317, 843
- Wang, N., Manchester, R. N., Zhang, J., Wu, X. J., Yusup, A., Lyne, A. G., Cheng, K. S., & Chen, M. Z. 2001, *MNRAS*, 328, 855
- Weltevrede, P., et al. 2010, *PASA*, 27, 64
- Yuan, J. P., Wang, N., Manchester, R. N., & Liu, Z. Y. 2010, *MNRAS*, 404, 289
- Zhou, A. Z., Xu, R. X., Wu, X. J., & Wang, N. 2004, *Astroparticle Phys.*, 22, 73




Fig. 1.— Glitch of PSR B2334+61: (a) variations of rotational frequency $\Delta\nu$ relative to the pre-glitch solution, (b) an expanded plot of $\Delta\nu$ where the mean post-glitch value has been subtracted from the post-glitch data, and (c) variations of the frequency first derivative $\dot{\nu}$. The vertical dashed line marks the glitch epoch. The plotted points represent fits of ν and $\dot{\nu}$ to 5 – 10 adjacent TOAs, with $\Delta\nu$ being the difference between the fitted value of ν and the value from the pre-glitch solution (extrapolated after the glitch).




Fig. 2.— Timing residuals for PSR B2334+61 relative to the pre-glitch solution (Table 1) for data up to one year after the glitch. For the upper plot just one exponential decay term was fitted, whereas for the lower plot two exponential decay terms were fitted. For further details, see text. The vertical dash line indicates the assumed epoch of glitch.




Fig. 3.— Timing residuals for the entire seven-year dataspan after fitting for two exponential recoveries and a permanent changes in $\dot{\nu}$ and $\ddot{\nu}$. The pulsar position and frequency parameters were held fixed at values obtained from the pre-glitch solution.

Table 1. Pre- and post-glitch timing parameters for PSR B2334+61

Parameter	Pre-glitch	Post-glitch
Right ascension (J2000)	23 ^h 37 ^m 05.762(10)	
Declination (J2000)	+61°51′01″53(7)	
Proper motion in RA dirn (mas yr ⁻¹)	−1 ^a	
Proper motion in Dec. (mas yr ⁻¹)	−15 ^a	
Epoch of position (MJD)	53100	
Frequency, ν (Hz)	2.01874888482(2)	2.01869363589(2)
Frequency derivative, $\dot{\nu}$ (10 ⁻¹⁵ s ⁻²)	−780.7074(5)	−788.332(3)
Frequency 2nd derivative, $\ddot{\nu}$ (10 ⁻²⁴ s ⁻³)	3.17(6)	14.46(9)
Frequency 3rd derivative, $\dddot{\nu}$ (10 ⁻³⁰ s ⁻⁴)	...	−0.37(2)
Frequency epoch (MJD)	53100.0	54521.0
Data span (MJD)	52486–53609	53998–55045
DM (pc cm ⁻³)	58.410	58.410
No. of TOAs	98	100
RMS timing residual (ms)	0.91	2.07
Reduced χ^2	1.04	4.88

^aFrom Hobbs et al. (2004)

Table 2. The glitch parameters

Parameter	Value
Glitch epoch (MJD)	53615(6)
$\Delta\nu_g/\nu$ (10^{-6})	20.5794(12)
$\Delta\dot{\nu}_g/\dot{\nu}$	0.156(4)
$\Delta\dot{\nu}_p$ (10^{-15} s $^{-2}$)	−8.684(17)
$\Delta\ddot{\nu}_p$ (10^{-24} s $^{-3}$)	8.4(2)
$\Delta\nu_{d1}$ (μ Hz)	0.19(3)
τ_{d1} (d)	21.4(5)
$\Delta\nu_{d2}$ (μ Hz)	0.119(4)
τ_{d2} (d)	147(2)
$Q = (\Delta\nu_{d1} + \Delta\nu_{d2})/\Delta\nu_g$	0.00751(5)
Data span (MJD)	52486–55045
Rms timing residual (ms)	1.72
Reduced χ^2	3.32